

Optical surface modes in the presence of nonlinearity and disorder

M. I. Molina,^{1,2} N. Lazarides,^{3,4} and G. P. Tsironis^{3,4}

¹*Departamento de Física, Facultad de Ciencias, Universidad de Chile, Casilla 653, Santiago, Chile*

²*Center for Optics and Photonics, Casilla 4016, Concepcion, Chile*

³*Department of Physics, University of Crete, P.O. Box 2208, 71003 Heraklion, Greece*

⁴*Institute of Electronic Structure and Laser, Foundation for Research and Technology–Hellas, P.O. Box 1527, 71110 Heraklion, Greece*

(Received 14 November 2011; published 31 January 2012)

We investigate numerically the effect of the competition of disorder, nonlinearity, and boundaries on the Anderson localization of light waves in finite-size, one-dimensional waveguide arrays. Using the discrete Anderson–nonlinear Schrödinger equation, the propagation of the mode amplitudes up to some finite distance is monitored. The analysis is based on the calculated localization length and the participation number, two standard measures for the statistical description of Anderson localization. For relatively weak disorder and nonlinearity, a higher disorder strength is required to achieve the same degree of localization at the edge than in the interior of the array, in agreement with recent experimental observations in the linear regime. However, for relatively strong disorder and/or nonlinearity, this behavior is reversed and it is now easier to localize an excitation at the edge than in the interior.

DOI: [10.1103/PhysRevE.85.017601](https://doi.org/10.1103/PhysRevE.85.017601)

PACS number(s): 05.45.Yv, 42.25.Dd, 42.65.Wi, 42.79.Gn

I. INTRODUCTION

A fundamental question concerning systems which are both disordered and nonlinear is whether or not Anderson localization [1] is weakened by the presence of nonlinearity. While it was originally developed in order to understand electronic transport in nonperiodic (disordered) solids, the concept of Anderson localization was later generalized to the localization of classical waves in disorder media [2]. The interaction of propagating waves and, in particular, of electromagnetic waves, when both disorder and nonlinearity are present can significantly affect localization and other phenomena [3].

Despite many efforts, that question has not been conclusively answered [4–11]. It thus seems that the answer depends on the relative strength of disorder and nonlinearity. For large nonlinearity, time-periodic and exponentially localized excitations in the form of discrete breathers may be generated, due to the self-trapping effect [12]. For small disorder strength, the discrete breathers are modulated to become localized modes [13]. The above theoretical results were accompanied by a series of experimental demonstrations of Anderson localization in optics [14] and Bose-Einstein condensates [15].

It was recently observed experimentally that Anderson localization in finite segments of disordered waveguide arrays in the linear regime is actually site dependent [16]. Specifically, a higher disorder strength is required to achieve the same degree of localization at the edge as in the interior (i.e., the “bulk”) of the array [16]. Here we are interested in the effect of the interplay of disorder and nonlinearity on the site dependence of wave-packet localization in one-dimensional (1D), disordered, finite-size arrays of coupled Kerr-type waveguides. Using the discrete nonlinear Schrödinger (DNLS) equation with diagonal (on-site) disorder, frequently referred to as the discrete Anderson–nonlinear Schrödinger (DANLS) equation, we calculate standard measures of Anderson localization in order to analyze the site dependence of the degree of localization for a wide range of nonlinearity and disorder strengths.

II. MODEL EQUATIONS AND STATISTICAL MEASURES

Consider a 1D array of N single-mode optical waveguides. In the framework of the coupled-mode formalism, the electric field $\mathcal{E}(x, z)$ propagating along the waveguides can be expanded as a superposition of the waveguide modes, $\mathcal{E}(x, z) = \sum_n E_n(z)\phi(x - n)$, where E_n is the complex amplitude of the single guide mode $\phi(x)$ centered at the n th site. The evolution equations for the modal amplitudes E_n are

$$i \frac{dE_n}{dz} + k_n E_n + V_{n,n-1} E_{n-1} + V_{n,n+1} E_{n+1} + \gamma |E_n|^2 E_n = 0, \quad (1)$$

where $n = 1, 2, 3, \dots, N$, E_n is the electric field [in units of $(W)^{1/2}$] on waveguide n at distance z (in meters), k_n is the propagation constant of guide n (in units of 1/m), $V_{n,m}$ is the coupling coefficient (in units of 1/m) between guides n and m , and γ [in units of 1/(W m)] is the nonlinear coefficient, defined by $\gamma = \omega_0 n_2 / c A_{\text{eff}}$, where ω_0 is the frequency of light, n_2 is the nonlinear refractive index of the guide, and A_{eff} is the effective area of the linear modes. Typical values in laser-written waveguide arrays are $N = 31$ guides, $\omega_0 = 1.326 \times 10^{-14}$ (1/s), and a propagation distance of 100 mm. The values of k_n can be adjusted by changing the (linear) refraction index of the guide. This is accomplished by varying the power or the writing speed of the laser beam that creates the waveguide array [16].

Equation (1) describes very well recent experiments in 1D disordered waveguide lattices and, moreover, it serves as a paradigmatic model for a wide class of physical problems where both disorder and nonlinearity are important. Disorder is introduced into the optical lattice by randomly choosing the propagation constants k_n from a uniform, zero-mean distribution. As a result, the lattice remains periodic on average and, to a very good approximation, the parameters $V_{n,n\pm 1}$ become independent of the site number n , i.e., $V_{n,n\pm 1} = V$. We render Eq. (1) dimensionless by defining $s = Vz$, $C_n =$

E_n/\sqrt{p} , $\chi = (\gamma p/V)^2$, and $\epsilon_n = k_n/V$, where p is the input beam power, $p = \sum_n |E_n|^2$. Equation (1) is then rewritten as

$$i \frac{dC_n}{dz} + \epsilon_n C_n + C_{n-1} + C_{n+1} + \chi |C_n|^2 C_n = 0, \quad (2)$$

where ϵ_n is a random number taken from the uniform distribution $[-\Delta, +\Delta]$. In current experiments, ranges of $[0, 2]$ for ϵ_n and $[0, 5]$ for χ are readily accessible [17].

In order to take into account the termination of the structure, we impose free boundary conditions at the edges, i.e., $C_0 = C_{N+1} = 0$. For $\chi \rightarrow 0$, Eq. (2) reduces to the original Anderson model while in the absence of disorder ($\epsilon_n = 0$), it reduces to the 1D DNLS equation [18] that is generally nonintegrable, and it conserves the norm $\sum_{n=1}^N |C_n|^2 = 1$ and the Hamiltonian \mathcal{H} .

To investigate the simultaneous interplay of disorder, nonlinearity, and boundary effects, initially we place a single-site excitation $C_n = \delta_{n,n_0}$ near, or at the boundary of the array. For a quantitative analysis we utilize two of the standard measures used in the description of Anderson localization; the participation number $P = \{\sum_{n=1}^N |C_n|^4\}^{-1}$ and the localization length ℓ , defined as the width of the envelope containing the localized profile. The participation number gives a rough estimate of the number of sites where the wave packet has significant amplitudes, and it is a useful measure for ascertaining localization effects in the case of partial localization. In this case, P will saturate at a finite value, indicating the formation of a localized wave packet.

III. STATISTICAL ANALYSIS

In the following, we set $V = 1$, while the nonlinearity parameter χ varies between 0 and 10, and the disorder width Δ takes on several different values. Since Anderson localization is essentially a statistical phenomenon, many realizations of disorder are needed to obtain meaningful averages for the quantities of interest. This is particularly true for low-dimensional systems. We typically use $n_R = 1000$ realizations in each run. The array contains $N = 200$ waveguides, and the maximum evolution “time” is $z = 100$ (except otherwise stated). In optics, nonlinearity is varied by changing the power content of the input beam. However, this is formally equivalent to keeping the norm of the wave packet fixed, and to varying the nonlinearity parameter χ . Equation (2) is integrated with a standard fourth-order Runge-Kutta algorithm with fixed time step. We compute the absolute squared profiles $\langle |C_n|^2 \rangle$, where the brackets denote averaging over all realizations n_R , hereafter referred to as Anderson mode profiles. Assuming that the Anderson modes have a z dependence which is a simple exponential function of the form $\langle |C_n|^2 \rangle = C_{\max}^2 e^{-z/\ell}$, the localization length ℓ can be computed via a χ^2 fitting procedure, with C_{\max}^2 being the numerically obtained maximum of $\langle |C_n|^2 \rangle$.

In Figs. 1–3 the inverse localization length ℓ^{-1} is shown as a function of χ for three different values of disorder strength. In all the cases displayed in these figures the initial wave packet is a single-site excitation placed at $n = n_0$, with $n_0 = 1$ (right at the edge), 2, 3, 5, and 10. In Fig. 1 (where $\Delta = 0.6$) we easily identify two different χ regimes; the weak and the strong non-




FIG. 1. (Color online) Inverse localization length ℓ^{-1} as a function of the nonlinearity strength χ . A single-site excitation is launched from $n_0 = 1$ (solid black), 2 (red dotted), 3 (green dashed), 5 (blue long-dashed), and 10 (indigo dotted-dashed), for a system with $N = 200$, $V = 1$, $z = 100$, $n_R = 1000$, and $\Delta = 0.6$.

linearity regimes, where ℓ^{-1} is small and large, respectively. That generally implies a lower degree of Anderson localization in the weak nonlinearity regime compared to that in the strong nonlinearity regime. The large ℓ^{-1} in the interval of χ values where all the curves fall one onto the other, indicates the existence of a highly localized mode due to the self-trapping effect. The characteristic nonlinearity strength, χ_c , that roughly distinguishes between the two regimes is the critical one for self-trapping to occur in the 1D DNLS equation [12]. In the weak nonlinearity regime another important feature appears; as can be seen in the figure, the $\ell^{-1}(\chi)$ curve obtained for excitations initially placed at the edge ($n_0 = 1$) is well below all the others (for which $n_0 > 1$). Thus, an excitation initially placed at the edge ($n_0 = 1$) leads to final wave packets that are less localized than those which have been initialized below the “surface” ($n_0 > 1$). This effect can be understood as the “repulsive” action of the boundary, reported in a previous work for surface modes in nonlinear periodic lattices [7], and it is in agreement with the experimental observations of Ref. [16].

As the disorder strength Δ is increased from 0.6 to 1.0 (Fig. 2), all the curves become flatter without showing any qualitative difference from those of Fig. 1. When Δ is increased to 2.0, however, we do observe qualitative differences (Fig. 3). For weak nonlinearity ($\chi \lesssim 4 \sim \chi_c$) there are no significant differences in the degree of localization for the Anderson modes resulting from initial excitations either at the edge or in




FIG. 2. (Color online) Inverse localization length ℓ^{-1} and localization length ℓ (inset) as a function of the nonlinearity strength χ . Same parameters and notation as in Fig. 1, but $\Delta = 1.0$.




FIG. 3. (Color online) Inverse localization length ℓ^{-1} and localization length ℓ (inset) as a function of the nonlinearity strength χ . Same parameters and notation as in Fig. 1, but $\Delta = 2.0$.

the interior of the array. Thus, it is as easy to localize a wave packet at the edge as it is in the interior in this case. However, for intermediate nonlinearities (from $\chi \gtrsim 4$ to $\chi \simeq 8$) it is more favorable to localize a wave packet at the edge than in the bulk, whereas for large nonlinearities $\chi \gtrsim 8$ it is as easy to localize a wave packet at the edge as it is in the bulk. Thus, for relatively strong disorder we observe a behavior that is strikingly different to what is observed in Figs. 1 and 2. The two different behaviors can be seen even more clearly by comparison of the localization length ℓ as a function of χ for $\Delta = 1$ and 2, shown in the insets of Figs. 2 and 3, respectively. Thus, the presence of strong disorder is capable of overcoming the repulsive character of the boundary for any value of χ and, moreover, it favors wave-packet localization at the edges for intermediate nonlinearities.

Typical examples of localized mode profiles both at, or close to the edge and the “bulk” are shown in Fig. 4, where the Anderson mode profiles $\langle |C_n|^2 \rangle$ are shown as a function of n for $\Delta = 0.6$. (Note that in this figure $z = 1000$.) The edge-




FIG. 4. Anderson mode profiles $\langle |C_n|^2 \rangle$ as a function of the site number n for $N = 200$, $V = 1$, $\Delta = 0.6$, $z = 1000$, $n_R = 1000$, and $\chi = 1$ [(a), (d)], $\chi = 2$ [(b), (e)], and $\chi = 3$ [(c), (f)]. Left panels denote the surface mode case ($n_0 = 1$) while right panels refer to the bulk mode case ($n_0 = 100$). Only part of the array sites are shown for clarity.




FIG. 5. (Color online) Averaged absolute squared profiles $\langle |C_n|^2 \rangle$ as a function of the site number n for $N = 200$, $V = 1$, $\Delta = 0.4$, $n_R = 1000$, $\chi = 2.5$, and $z = 100$. These profiles result from a single-site initial excitation at $n_0 = 1$ (black solid), $n_0 = 2$ (red dotted), and $n_0 = 3$ (green dashed). Only part of the array sites are shown for clarity.

localized modes are significantly more extended than the bulk modes, even though the former are not all localized exactly at the edge. This is because of the small disorder strength Δ , which allows the repulsive force of the boundary on the mode to dominate and move the mode maximum slightly toward the bulk. However, similar profiles (not shown) are also obtained for $\Delta = 1.0$, that is large enough to keep the localized modes at their initial location.

Moreover, single-site excitations initialized at different sites n_0 can be pushed by the boundary toward the interior and form Anderson modes at the same final site. These modes are different, at least for finite propagation distances z ; they differ in the degree of localization, leading to a multiplicity of Anderson modes having their maximum at the same site of the lattice (Fig. 5). For the particular value of χ used for Fig. 5, three single-site excitations initialized at different n_0 have formed (after they have been propagated up to $z = 100$) three distinct Anderson localized modes whose maximum is located at the same lattice site (i.e., at $n = 3$).

Finally, let us look at the participation number P as a function of z for wave packets that are initially localized at the edge ($n_0 = 1$) and in the bulk ($n_0 = 100$) of the waveguide array. The logarithmic plots are shown in Figs. 6 and 7, respectively, for several values of χ and weak disorder. Comparing the curves in these figures corresponding to the




FIG. 6. (Color online) Logarithm of the averaged participation number $\log_{10}(\langle P \rangle)$ as a function of $\log_{10}(z)$ for the surface case ($n_0 = 1$) and $N = 200$, $V = 1$, $\Delta = 0.6$, $n_R = 1000$, and $z = 1000$. The values of χ are shown in the figure.




FIG. 7. (Color online) Logarithm of the averaged participation number $\log_{10}(\langle P \rangle)$ as a function of $\log_{10}(z)$ for the bulk case ($n_0 = 100$) and $N = 200$, $V = 1$, $\Delta = 0.6$, $n_R = 1000$, $z = 1000$, and $n_0 = 100$. The values of χ are shown in the figure.

same χ , we see that those for $n_0 = 100$ are shifted to higher P values than those for $n_0 = 1$. Thus, single-site excitations initialized at $n_0 = 100$ have, while propagating along z , a larger number of sites where the wave packet has significant amplitudes. However, the excitations initialized at $n_0 = 100$ exhibit a higher degree of localization than those initialized at $n_0 = 1$ (see also Fig. 4). It is also interesting to see how the curves in each figure change as a function of χ . For $\chi = 6$, well above χ_c , we see that the wave packet remains localized at a single site, independently of n_0 . For $\chi = 4 \sim \chi_c$, the $\log_{10}(\langle P \rangle)$ vs $\log_{10}(z)$ curve increases slowly with increasing $\log_{10}(z)$, and it saturates at a finite value around $z \simeq 100$, indicating the formation of a wave packet highly localized around n_0 . For the nonlinearity strengths that are less than χ_c , the $\log_{10}(\langle P \rangle)$ vs $\log_{10}(z)$ curves exhibit qualitatively the same behavior. There is an increase with increasing $\log_{10}(z)$ which is slowed down after some z specific to each χ value, and indicates significant delocalization of the initially single-site wave packet. Delocalization is stronger for decreasing nonlinearity strength. However, those curves do not seem to

saturate, implying that the corresponding Anderson localized modes may delocalize further at longer propagation distances.

IV. CONCLUDING REMARKS

We have performed extensive calculations with the 1D DANLS equation in order to clarify some aspects of the interplay between boundary effects, disorder, and nonlinearity, in finite-size waveguide arrays. In particular, we attempt to clarify the site dependence of Anderson localization that results from that interplay. We computed two standard measures of localization for discrete systems for varying nonlinearity and disorder strengths, and we observed two strikingly different behaviors depending on the strength of the disorder. For weak to moderate disorder, we distinguish two different nonlinearity regimes: weak and strong, for values of χ roughly below and above χ_c , respectively. In the weak nonlinearity regime it is easier to localize a wave packet in the interior of the array than at the edge, which is in agreement with the experiments in the linear regime [16]. In the strong nonlinearity regime it is as easy to localize a wave packet at the edge as it is in the interior. However, for relatively strong disorder, this behavior is reversed, at least for intermediate nonlinearities, and it is now easier to localize a wave packet at the edge than in the bulk. For weak and very strong nonlinearities there is no significant site dependence on the degree of wave-packet localization. The results presented here obviously hold for finite propagation distance z , an important case of practical interest for experimentalists.

ACKNOWLEDGMENTS

M.I.M. acknowledges support from Fondecyt Grant No. 1080374 and Programa de Financiamiento Basal de Conicyt (Grant No. FB0824/2008). G.P.T. and N.L. acknowledge partial support by the THALES Project MACOMSYS, funded by the ESPA Program of the Ministry of Education of Greece.

-
- [1] P. W. Anderson, *Phys. Rev.* **109**, 1492 (1958).
 [2] S. John, *Phys. Rev. Lett.* **58**, 2486 (1987).
 [3] *Disorder and Nonlinearity*, edited by A. R. Bishop, D. K. Campbell, and St. Pnevmatikos (Springer-Verlag, Berlin, 1989); *Nonlinearity with Disorder*, edited by F. Kh. Abdullaev, A. R. Bishop, and St. Pnevmatikos (Springer-Verlag, Berlin, 1991).
 [4] M. I. Molina, *Phys. Rev. B* **58**, 12547 (1998).
 [5] G. Kopidakis and S. Aubry, *Phys. Rev. Lett.* **84**, 3236 (2000).
 [6] G. Kopidakis, S. Komineas, S. Flach, and S. Aubry, *Phys. Rev. Lett.* **100**, 084103 (2008).
 [7] M. I. Molina, R. A. Vicencio, and Y. S. Kivshar, *Opt. Lett.* **31**, 1693 (2006).
 [8] Yu. S. Kivshar, S. A. Gredeskul, A. Sanchez, and L. Vazquez, *Phys. Rev. Lett.* **64**, 1693 (1990).
 [9] A. S. Pikovsky and D. L. Shepelyansky, *Phys. Rev. Lett.* **100**, 094101 (2008).
 [10] S. Flach, D. O. Krimer, and Ch. Skokos, *Phys. Rev. Lett.* **102**, 024101 (2009).
 [11] S. Fishman, Y. Krivolapov, and A. Soffer, e-print arXiv:1108.2956v1.
 [12] M. I. Molina and G. P. Tsironis, *Physica D* **65**, 267 (1993); *Int. J. Mod. Phys. B* **9**, 1899 (1995).
 [13] M. V. Ivanchenko, *Phys. Rev. Lett.* **102**, 175507 (2009).
 [14] T. Pertsch *et al.*, *Phys. Rev. Lett.* **93**, 053901 (2004); T. Schwartz *et al.*, *Nature (London)* **446**, 52 (2007); Y. Lahini *et al.*, *Phys. Rev. Lett.* **100**, 013906 (2008).
 [15] J. Billy *et al.*, *Nature (London)* **453**, 891 (2008); G. Roati *et al.*, *ibid.* **453**, 895 (2008).
 [16] A. Szameit *et al.*, *Opt. Lett.* **35**, 1172 (2010).
 [17] A. Szameit (private communication).
 [18] P. G. Kevrekidis, *The Discrete Nonlinear Schrödinger Equation* (Springer-Verlag, Berlin, 2009).